arXiv:hep-ph/0402099v1 10 Feb 2004

ONE-LOOP CONTRIBUTIONS OF CHARGINOS AND NEUTRALINOS TO W^+W^- PRODUCTION IN E^+E^- COLLISIONS

KAORU HAGIWARA
Theory group, KEK, Tsukuba, Ibaraki, 305-0851,Japan
E-mail: kaoru.hagiwara@kek.jp

SHINYA KANEMURA
Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan
E-mail: kanemu@het.phys.sci.osaka-u.ac.jp

YOSHIAKI UMEDA*
Institute of physics, National Chiao Tung University, Hsinchu 300, Taiwan
E-mail: umeda@faculty.nctu.edu.tw

We study one-loop effects of charginos and neutralinos on the helicity amplitudes for the process $e^+e^- \to W^+W^-$ in the MSSM. The calculation is tested by using two methods. First, the sum rules among form factors which result from BRS invariance. Second, the decoupling property in the large mass limit is used to test the overall normalization of the amplitudes. The one-loop contribution to the helicity amplitudes is shown as a function of μ Higgs parameter and $\sqrt{s}$. We also study the effects of the CP-violating phase.

1. Introduction

We discuss the one-loop charginos and neutralinos contributions to the $e^+(k,\tau)e^-(\overline{k},\overline{\tau}) \to W^+(p,\lambda)W^-(\overline{k},\overline{\lambda})$ in the MSSM[1]. Here k and p are the momentum, and τ and λ denote helicities. The mass matrix of charginos or neutralinos are express by 2×2 or 4×4 matrix, respectively. The helicity amplitudes can be written by using 16 basis tensors $T_i^{\mu\alpha\beta}$

$$M_\tau^{\lambda\overline{\lambda}} = \sum_{i=1}^{16} F_{i,\tau}(s,t)\, j_\mu(k,\overline{k},\tau) T_i^{\mu\alpha\beta} \epsilon_\alpha(p,\lambda)^* \epsilon_\beta(\overline{p},\overline{\lambda})^* \ , \tag{1}$$

---

*Supported by BMBF, contract nr. 05 HT1PAA 4 and NSC 92-2811-M-009-029.

Figure 1. One-loop contributions to M_τ^{00} helicity amplitudes are shown.



where j_μ is the electron current and ϵ_α is the polarization vectors for W. All the dynamical information is contained in the form factors $F_{i,\tau}(s,t)$. Among the 16 form factors, physically polarized W boson are described by nine form factors. To test the physical form factors of $F_{i,\tau}(s,t)$(i=1−9), we calculate unphysical form factors of $F_{i,\tau}(s,t)$(i=10−13) together with the form factors of $H_{i,\tau}(s,t)$(i=1−4) of the helicity amplitudes of $e^+(k,\tau)e^-(\overline{k},\overline{\tau}) \to \omega^+(p)W^-(\overline{k},\overline{\lambda})$ (ω^+: Nambu-Goldstone boson), by using basis tensors $S_i^{\mu\alpha}$

$$M_\tau^\lambda(e^+e^- \to \omega^+W^-) = i\sum_{i=1}^{4} H_{i,\tau}(s,t)\, j_\mu(k,\overline{k},\tau) S_i^{\mu\alpha} \epsilon_\alpha(p,\lambda)^* \, . \tag{2}$$

2. Test

In order to ensure the correctness of our calculation, we examine the BRS invariance of one-loop helicity amplitudes and the decoupling behavior of the SUSY effects in the large mass limit of charginos and neutralinos.

2.1. *The BRS invariance*

The linear relation between $e^+e^- \to W_S^+W_P^-$ and $e^+e^- \to \omega^+W_P^-$ helicity amplitudes is obtained from BRS invariance, where index P and S denote physical and scalar polarization, respectively. We obtain three sum rules among the form factors of $\widetilde{F}_{i,\tau}$ and $\widetilde{H}_{3,\tau}$ from BRS invariance[2]

$$\sum_{j=1}^{16} \xi_{i,j} F_{j,\tau}(s,t) = C_{mod}^{BRS} H_{i,\tau}(s,t)\, , \tag{3}$$

where i runs from 1 to 3 and $\xi_{i,j}$ are determined by kinematical parameters. For the factor $C_{\rm mod}^{BRS}$, see Ref. [2]. We numerically confirmed that the Eq. (3)

Figure 2. The effects of CP-violating phases are shown.



Table 1. Input parameters and mass spectra. The $\tan\beta = 3$ for all cases.

	A	B	C	D	E
Parameter					
μ (GeV)	+120	+145	+400	+1000	+130
M_2 (GeV)	541	242	125	115	158
φ_1	0	0	0	0	$\frac{2}{3}\pi$
φ_μ	0	0	0	0	$\frac{2}{3}\pi$
Mass spectra (GeV)					
$m_{\widetilde{\chi}_1^-}$	110	110	110	110	110
$m_{\widetilde{\chi}_2^-}$	555	283	420	1007	207
$m_{\widetilde{\chi}_1^0}$	99	81	60	57	75
$m_{\widetilde{\chi}_2^0}$	123	150	111	110	105
$m_{\widetilde{\chi}_3^0}$	285	150	403	1002	154
$m_{\widetilde{\chi}_4^0}$	555	285	422	1007	205

agrees more than 11 digit numerically.

2.2. *Decoupling limit*

The one-loop effects of the SUSY particles should decouple from the low energy observable in the large mass limit. By using this fact, we can test the overall normalization of the amplitudes. We check that the differential cross section of the MSSM become that of the SM in the large mass limit of the gaugino mass M_1 and M_2, and μ higgs term. See Fig. 1 of Ref. [1].

3. Numerical results

We present the one-loop effects to the helicity amplitudes. We assume the GUT relation $M_1 = 5M_2\hat{s}^2/3\hat{c}^2$, light chargino mass $m_{\widetilde{\chi}_1^-} = 110$GeV, and scattering angle $\theta = 90°$.

3.1. *The chargino and neutralino contribution to M_τ^{00}*

Among the helicity amplitudes, the loop correction is the largest in 00 helicity amplitude. In Fig. 1(a), we show the μ dependence for $\tan\beta = 3$ or 50. The loop effects are about 0.7% at $|\mu| = 1000$GeV for $\tau = -1$, whereas they remain small for $\tau = +1$ at around 0.1% level. In Fig. 1(b), the $\sqrt{s}$ dependence for four sets of parameter (sets A to D of Table 1) is shown. Sharp peak can be seen for each curve, which correspond to the thresholds of pair production of the lightest chargino at $\sqrt{s} = 220$GeV for $\tau = -1$. The corrections are negative for $\tau = +1$.

3.2. *CP-violating effects*

There are new CP-violating phases in the MSSM. In the chargino and neutralino sectors, it arises from the μ Higgs mass term $\mu e^{i\varphi_\mu}$ and the gaugino mass parameters $M_1 e^{i\varphi_1}$ for the most general case when we take the phase of M_2 be 0 by rephasing. Large CP-violating phases in chargino and neutralino sectors are possible without contradicting the EDM constraints of electrons and neutrons, if the parameters for sleptons and squarks are adjusted. In Figs. 2, the real part (solid curve) and the imaginary part (dotted curve) of f_4^Z, f_6^γ, and f_6^Z of Ref. [1,2] are shown as a function of $\sqrt{s}$ for the parameters of set E in Table 1. These form factors take their maximum or minimum at around $\varphi_\mu = \varphi_1 = 2/3\pi$ or $4/3\pi$. We can measure the CP-violating effects of f_4 and f_6 by the difference of helicity amplitudes $M^{\pm 0}$ and $M^{\mp 0}$. The CP-violating effects in the helicity amplitudes can be of the order (0.1%).

4. Conclusion

The correction due to chargino and neutralino contributions to $e^+e^- \to W^+W^-$ helicity amplitudes can be of the order (1%) and the CP-violating effects can be of the order (0.1%).

References

1. K. Hagiwara, S. Kanemura, M. Klasen and Y. Umeda, *Phys. Rev.* **D68**, 113011 (2003).
2. S. Alam, K. Hagiwara, S. Kanemura, R. Szalapski and Y. Umeda, *Phys. Rev.* **D62**, 095011 (2000); S. Alam, K. Hagiwara, S. Kanemura, R. Szalapski and Y. Umeda, *Nucl. Phys.* **B541** 50 (1999); K. Hagiwara, S. Kanemura and Y. Umeda, *Proceedings of 30th International Conference on High-Energy Physics* (ICHEP 2000),[hep-ph/0009200].